

2000% More Fetal Deaths Occurred After The C-19 Rollout! CDC Still Recommends Vaccination For Pregnant Women!

Description

USA: The FDA and CDC's criminal actions keep mRNA vaccines in the market, even though there is too much evidence showing they are harmful.

More than one million cases of deaths and injuries have been reported. The US Government VAERS started to be reported in the corporate pro-Pharma media, as insurance companies continued to report on their data from 2021 when the vaccines were rolled out, showing a dramatic increase in life insurance and health insurance claims due to C-19 vaccines.

Join The True Defender Telegram Chanel Here: https://t.me/TheTrueDefender

Check this out:

68,000% Increase in Strokes as FDA and NIH Secretly Study Reports of Neurological Injuries After COVID-19 Vaccines

German Health Insurance Claims Show 31,254 Deaths Following COVID-19 Vaccines While Official Government Stats Report Only 2,255


CDC Withholding Data on COVID-19 Vaccines Because "They might be misinterpreted as the vaccines being ineffective" and "The figures would be misrepresented by anti-vaccine groups"

Youngsters suffer from heart diseases after the C-19 vaccination, and the CDC and FDA confessed that myocarditis and pericarditis are possible side effects of the vaccine.

One fact is shown below. The shots kill unborn children, and the CDC hasn't updated their recommendations for C-19 vaccines and pregnant women despite this damaging data.


From the 2/18/2022 release of VAERS data:

Found 3,725 cases where Vaccine is COVID19 and Symptom is Aborted pregnancy or Abortion or Abortion complete or Abortion complete or Abortion induced or Abortion induced or Abortion induced complete or Abortion induced incomplete or Abortion late or Abortion missed or Abortion of ectopic pregnancy or Abortion spontaneous or Abortion spontaneous complete or Abortion spontaneous incomplete or Ectopic pregnancy or Ectopic pregnancy termination or Ectopic pregnancy with contraceptive device or Foetal cardiac arrest or Foetal death or Premature baby death or Premature delivery or Ruptured ectopic pregnancy or Stillbirth



From the 2/18/2022 release of VAERS data:

Found 3,725 cases where Vaccine is COVID19 and Symptom is Aborted pregnancy or Abortion or Abortion complete or Abortion complete or Abortion induced or Abortion induced complete or Abortion induced incomplete or Abortion late or Abortion missed or Abortion of ectopic pregnancy or Abortion spontaneous or Abortion spontaneous complete or Abortion spontaneous incomplete or Ectopic pregnancy or Ectopic pregnancy termination or Ectopic pregnancy with contraceptive device or Foetal cardiac arrest or Foetal death or Premature baby death or Premature delivery or Ruptured ectopic pregnancy or Stillbirth



The latest VAERS data shows that there have been 3.725 fetal deaths after the C-19 vaccinations.

Most of the deaths have occurred due to the Pfizer vaccine.

The CDC shared that there have been 550million doses of C-19 vaccines administered only in the US.

It means that for every 147.651 doses of C-19 vaccines, there has been one fetal death.

There were more than 4 billion doses of the vaccines administered in the US from 2006 to 2019. Throughout this period, 1.369 fetal deaths have been reported in VAERS. Most deaths have been reported from the HPV vaccines.

It means that one death has been reported to VAERS from 2006 to 2019 for every 2.989,596 doses administered.

It's a 1.925 increase in fetal deaths after the rollout of the C-19 vaccine compared to all vaccines administered during the 14 years from 2006 to 2019.

CDC shared:

"COVID-19 vaccination is recommended for people who are trying to get pregnant now or might become pregnant in the future, as well as their partners. In addition, everyone who is pregnant, breastfeeding, trying to get pregnant now, or might become pregnant in the future, should get a booster shot. CDC recommends that people who are starting their vaccine series or getting a booster dose get either Pfizer-BioNTech or Moderna (mRNA COVID-19 vaccines), but the J&J/Janssen COVID-19 vaccine may be considered in some situations.

There is no recommendation for routine pregnancy testing before receiving a COVID-19 vaccine. If you are trying to become pregnant or may want to become pregnant, you do not need to avoid receiving a COVID-19 vaccine. Those who are trying to become pregnant do not need to avoid pregnancy after the COVID-19 vaccination. If you get pregnant after receiving your first shot of a COVID-19 vaccine that requires two doses (i.e., Pfizer-BioNTech COVID-19 vaccine or Moderna COVID-19 vaccine), you should get your second shot to get as much protection as possible. (Source.)"

Watch the videos below:

by Addison Wilson

Category

- 1. Health-Wellness-Healing-Nutrition & Fitness
- 2. Main
- 3. Politics-Geopolitics-Gov.-Events

Date Created

03/04/2022