

Billions of Dangerous Spike DNA's Molecules inside Covid mRNA Vaccines

Description

In the cover image an image of the research of Dr. Kevin McKernan and the molecular biologist Jessica Rose who confirmed its importance

The scientific revelations arriving from the United States of America not only represent yet another disturbing alarm on the dangerousness of gene serums against SARS-Cov-2 but could be able to explain many still mysterious phenomena such as Covid break-through, i.e. when the Covid-19 "holes" the vaccine protection by triggering the pathology in a serious or even lethal symptomatic way, and like Long Covid.

The alarm about the enormous number of Spike's DNA molecules circulating in the human body was launched as early as summer 2021 by US Air Force lieutenant colonel and surgeon Theresa Long when she wrote an affidavit to justify her decision not to want to fly vaccinated pilots at risk of heart attacks from myocarditis.

Alarming Contamination of mRNA vaccine

The research of genomics expert Dr. Kevin McKernan was reported by Will Jones on the American website The Daily Skeptic which was founded by journalists and academics in April 2020 under the name Lockdown Skeptic focusing precisely on the pandemic emergency and contains a striking discovery summarized in two lines.

"The contamination of mRNA vaccines with DNA is far greater than initially thought at up to 35%, and the DNA's role in inducing human cells to produce the spike protein long term has been confirmed, according to the latest research".

The study conducted in the biochemical laboratories of McKernan (founder of Medicinal Genomics) has currently only been published in the form of a scientific article on his Substack Anandamide and not in specialized medical journals.

But it is very reliable because it was supported by the analyses of the American biochemist and immunologist Jessica Rose,

author of many important studies on the dangers of mRNA vaccines that we have relaunched on Gospa News.

Furthermore, the discovery of the long permanence of the toxic protein Spike in the blood and in human organs has already been confirmed by other researches, including that of a group of Italian doctors.

Primarily from the study by Professor Markus Aldén of Lund University in Malmoe (Sweden) which sounded the alarm about possible DNA alterations caused by mRNA gene serums and induced the judge of the Court of Florence, Dr. Susanna Zanda, to forward his civil sentence to the Public Prosecutor's Office of Rome to investigate the damage and deaths from vaccines.

Dr Michael Mörz of the Institute of Pathology in Dresden, Germany, found traces of the Spike protein in the brain of a deceased trivaccinate.

The Toxic Spike of SARS-Cov-2 Enhanced in the Laboratory

The danger of this toxic protein was also reported by retired US Army lieutenant colonel Lawrence Sellin, an expert in microbiology by virtue of his work at Fort Detrick (Maryland), the most important American military bacteriological laboratory

Sellin explained in a scientific article how SARS-Cov-2 was artificially constructed by enhancing the Spike's viral load with the Gain-of-Function technique and alerted the scientific authorities to the fact that the same extremely dangerous virulence could also be reproduced in the mRNA and mDNA gene sera although this protein was genetically modified and attenuated only to trigger the immunological reaction.

His findings were recently confirmed by two studies published in PubMed, the official scientific archive of the National Institutes of Health (NHS) in the USA. The first by German and American scientists, the last, exclusively published by us, by Austrian and Canadian scientists.

The Permanence of Dangerous Plasmids in the Blood

Genomics expert McKernan today reveals the exact mechanism underlying the long permanence in the blood of the Spike inoculated with vaccines. He attributes it to the presence of small circular DNA molecules, plasmids, capable of reproducing between human and bacterial cells, thus explaining the bacteriophage nature of SARS-Cov-2 which makes it difficult to eliminate from the human body as it nestles in the Intestinal metabiota, as already highlighted by an Italian study.

It should be recalled that the first experiments conducted in the Wuhan Institute of Virology on laboratory-built chimeric SARS were the construction of a recombinant virus in which a plasmid infected with HIV was added to the SARS pathogen, probably in an attempt to research a vaccine antiAIDS as supported by Luc Montagnier who found HIV sequences inside SARS-Cov-2.

There is obviously no certainty that the Covid-19 virus was born precisely from those experiments then developed together with the University of North Carolina under the supervision of Anthony Fauci.

Therefore, the high contamination of Spike's DNA plasmids in vaccines could be useful in explaining many of the serious side effects.

But above all the neurocerebral disorders of the so-called Long Covid which the American National Institute of Health found almost identical to multiple neurological adverse reactions following the inoculation of gene serums.

In the same way, it supports the tremendous hypothesis that the Covid break-through are not determined by an ineffectiveness of the mRNA gene serums but that it is actually the vaccines that cause serious and lethal SARS-COV-2 pathologies as hypothesized by Gospa News after the alarming studies published in various parts of the world on this phenomenon which is minimized in Europe and even ignored in Italy.

by Carlo Domenico Cristofori

mRNA Vaccines Contain DNA That May Turn Human Cells Into Long-Term Spike Protein Factories

Will Jones - originally published by The Daily Sceptic

All links to Gospa News articles have been added aftermath in relation to the topics highlighted

Earlier this month, the *Daily Sceptic* reported on the work of Dr. Kevin McKernan and his team who had subjected the mRNA vaccines from Pfizer and Moderna to deep sequencing analysis and found alarming levels of DNA contaminants known as plasmids. These are small circular DNA molecules that in principle can self-replicate in bacterial and human cells and induce the cell to produce the SARS-CoV-2 spike protein long term. Each vaccine dose was found to contain billions of these plasmids.

The Moderna vaccine appeared to contain DNA contamination at around the 'safe' level set by the European Medicines Agency (EMA) at the equivalent of one part per 3,000 mRNA molecules, though it's not clear how safe this level really is. The Pfizer vaccine, on the other hand, was found to contain DNA contamination at 10 times the 'safe' level, at one plasmid per 350 mRNA molecules.

The DNA is part of the vaccine manufacturing process, providing the blueprint for the mRNA, but it should have been removed to at least the 'safe' level, though was not for reasons that are unclear.

DNA Molecules in Gene Serums 100 Times Higher than Safe Levels

Now, Dr. McKernan and his team have undertaken further analysis and found that the level of DNA contamination is much greater than originally reported, with up to 35% of the vaccine product being this DNA contamination. They write:

This equates to 20-35% of the nucleic acid in each vaccine being expression vector. This is several orders of magnitude over the the EMAs limit of 330ng/mg. With these levels of contamination, RT

activity from LINE-1 is not a prerequisite for genome integration.

Molecular biologist Dr. Jessica Rose explains that this means each dose may contain trillions of DNA molecules, 100 times greater than previously reported: "The left-over expression vectors used to manufacture the mRNAs are at contamination levels 100-fold higher than originally proposed and imply trillions of DNA molecules per dose. This has implications for integration into our genome."

The precise level of contamination is unknown as these are estimates with a wide margin of error. It may also vary by batch. What is certain, however, is that the contamination of both Pfizer and Moderna vaccines is way above any official 'safe' level.

In the original analysis, Dr. McKernan had been looking primarily for RNA contamination and had used an additive that can suppress DNA amplification. By looking specifically for DNA contamination he and his team found that the amount of DNA present was far greater than the initial technique had indicated.

Damage from Vaccines due to Plasmids

Further analysis by Dr. McKernan and his team has also confirmed that the plasmids are intact and capable of self-replicating, and that the relevant promoters are present that allow them to express mRNA for spike protein in human cells (and not just in bacteria).




Figure 3. Production of m1 Ψ mRNAs by *in vitro* transcription. Left: Components of *ir* reaction. Right: Incorporation of m1 Ψ -triphosphate into RNA is guided by m1 Ψ 's abil base pair with adenine of the DNA template in the T7 RNA polymerase active site.

The mechanism of self-replication of plasmids in human cells

This indicates that these DNA plasmids are likely to survive for long periods, be taken up by cellsinside the body and induce the cells to produce spike protein for an indefinite period of time.

It is thought this could explain the observed persistence of spike protein in the blood of vaccinated persons for weeks or months after injection, which is believed by experts to be a contributor to adverse effects of the COVID-19 vaccines.

Dr. Jessica Rose notes: "It is more than likely that these adverse effects are the direct result of the contamination illuminated by Kevin and his team."

These findings are obviously highly disturbing. Regulators ought to be making a priority of looking into these issues for themselves and, if confirmed, taking the appropriate action including removing the products from the market.

Will Jones – originally published by The Daily Sceptic

By Fabio G. C. Carisio **Category**

- 1. Crime-Justice-Terrorism-Corruption
- 2. Disasters-Crisis-Depopulation-Genocide
- 3. Health-Wellness-Healing-Nutrition & Fitness
- 4. Main
- 5. NWO-Deep State-Dictatorship-Tyrrany

Date Created

05/25/2023