

FDA had Data Showing 82% – 97% of Pregnant Women Injected with the Pfizer COVID-19 Vaccine Lost Their Babies Before Approving the Shots

Description

USA: Anne Reed of Operation Rescue [reported last week](#) that recently released documents from Pfizer that were submitted to the FDA prior to approving their COVID-19 vaccine in early 2021 showed that 82% – 97% of the documented pregnancy outcomes resulted in death in [their post-marketing analysis](#).

This is more evidence that Pfizer and the FDA colluded together to conceal damaging data related to the roll out of the Pfizer COVID-19 vaccines.

Back in May of 2021, Operation Rescue published a report from a whistleblower who had examined the Pfizer documents submitted to the European Medicines Agency (EMA) which authorized the Pfizer COVID-19 vaccines for emergency use in the UK back in December of 2020, and among those documents (which we assume were also submitted to the FDA but not yet made public) were the results of animal trials showing serious birth defects occurred in rat specimens injected with the vaccines. See:

[EMA Pfizer Documents on Experimental COVID-19 mRNA Shots Reveal Animal Studies were Conducted during Trials – Risks to Pregnancy being Concealed but Verified by VAERS Data](#)

When the EMA originally authorized the Pfizer COVID-19 vaccines, they included a warning about injecting child-bearing women:

Fertility, pregnancy and lactation

Pregnancy

There are no or limited amount of data from the use of COVID-19 mRNA Vaccine BNT162b2. Animal reproductive toxicity studies have not been completed. COVID-19 mRNA Vaccine BNT162b2 is not recommended during pregnancy.

For women of childbearing age, ***pregnancy should be excluded before vaccination***. In addition, women of childbearing age should be advised to avoid pregnancy for at least 2 months after their second dose.

Breast-feeding

It is unknown whether COVID-19 mRNA Vaccine BNT162b2 is excreted in human milk. A risk to the newborns/infants cannot be excluded. COVID-19 mRNA Vaccine BNT162b2 should not be used during breast-feeding.

Fertility

It is unknown whether COVID-19 mRNA Vaccine BNT162b2 has an impact on fertility. ([Source.](#))

However, when the FDA issued an EUA for the Pfizer COVID-19 vaccines shortly after the EMA, in December of 2020, they did NOT include any warnings for pregnant and child-bearing women. See:

Unlike UK, U.S. FDA Allows Pregnant and Nursing Women to Receive Experimental Pfizer COVID Vaccine

As of the 5/20/2022 release of VAERS data, there have been 4,202 cases recorded of fetal deaths following COVID-19 vaccines for the past 18 months since these vaccines were issued EUAs. ([Source.](#))

By way of contrast, performing the exact same search in VAERS for the previous 30 years (360 months) there were only 2,239 fetal deaths recorded following all FDA-approved vaccines during those 30+ years. ([Source.](#))

That's a 3,653% increase in dead babies following COVID-19 vaccines, compared to deaths of pre-born children following vaccines for the previous 30+ years.


COURT ORDERED RELEASE OF PFIZER DOCUMENT REVEALS 82% – 97% OF VACCINATED PREGNANT WOMEN LOST THEIR BABIES

by Anne Reed
[Operation Rescue](#)

A recently released confidential document reveals that a startling majority of in utero babies died after pregnant women were injected with the Pfizer vaccine.

Public Health and Medical Professionals for Transparency (PHMPT) requested documents related to vaccine licensing via a [Freedom of Information Act Request](#), then [sued the Food and Drug Administration \(FDA\)](#). In early January 2022, a federal judge ordered the FDA incrementally release the documents it claimed would take decades to process. PHMPT has posted [Pfizer's confidential documents](#) as they have become available.

In the most recent release, on May 2, 2022, a document titled '[reissue_5.3.6 postmarketing experience.pdf](#)' disclosed on page 12 that by February 8, 2021, 270 women had received the mRNA injection during pregnancy. But 238 cases were apparently not followed ("no outcome provided"). And, therefore, the pregnancy outcomes for those women are unknown.




The mere fact that 88% of the pregnant women injected were not followed throughout their pregnancies is deeply concerning since 124 of the 270 pregnant women had some type of adverse reaction (49 nonserious, 75 serious) according to page 12 of the same document.

Pregnancy cases: 274 cases including:

- 270 mother cases and 4 fetus/baby cases representing 270 unique pregnancies (the 4 fetus/baby cases were linked to 3 mother cases; 1 mother case involved twins).
- Pregnancy outcomes for the 270 pregnancies were reported as spontaneous abortion (23), outcome pending (5), premature birth with neonatal death, spontaneous abortion with intrauterine death (2 each), spontaneous abortion with neonatal death, and normal outcome (1 each). No outcome was provided for 238 pregnancies (note that 2 different outcomes were reported for each twin, and both were counted).
- 146 non-serious mother cases reported exposure to vaccine in utero without the occurrence of any clinical adverse event. The exposure PTs coded to the PTs Maternal exposure during pregnancy (111), Exposure during pregnancy (29) and Maternal exposure timing unspecified (6). Trimester of exposure was reported in 21 of these cases: 1st trimester (15 cases), 2nd trimester (7), and 3rd trimester (2).
- 124 mother cases, 49 non-serious and 75 serious, reported clinical events, which occurred in the vaccinated mothers. Pregnancy related events reported in these cases coded to the PTs Abortion spontaneous (25), Uterine contraction during pregnancy, Premature rupture of membranes, Abortion, Abortion missed, and Foetal death (1 each). Other clinical events which occurred in more than 5 cases coded to the PTs Headache (33), Vaccination site pain (24), Pain in extremity and Fatigue (22 each), Myalgia and Pyrexia (16 each), Chills (13) Nausea (12), Pain (11), Arthralgia (9), Lymphadenopathy and Drug ineffective (7 each), Chest pain, Dizziness and Asthenia (6 each), Malaise and COVID-19 (5 each). Trimester of exposure was reported in 22 of these cases: 1st trimester (19 cases), 2nd trimester (1 case), 3rd trimester (2 cases).

Among those 34 pregnancies that are known, the report indicates that 28 babies died either in utero or upon birth. Only one outcome was reported as normal, and the remaining five were reported as “pending.”



In analyzing Pfizer’s own data, it is clear that 82% – 97% of the documented pregnancy outcomes resulted in death. (The 15-point variation is dependent on the final outcome of those in the “outcome pending” category.)

“This report confirms information we released last year,” said Operation Rescue president Troy Newman. “We look forward to seeing some kind of accountability put in place for those responsible for hiding this crucial data from the public.”

See Operation Rescue’s previous coverage on pregnancy issues related to Covid-19 vaccines:

- [Whistleblower reveals pregnancy complications](#)
- [Whistleblowers and other experts warn of increased risk of infertility](#)
- [Bombshell interview with COVID 19 Task Force Whistleblower – Warns of Vaccine Risks to Pregnant Women](#)

**Note that two different outcomes were reported for twins, and both were counted.*

Read the full article at [Operation Rescue](#).

by **Brian Shilhavy**

Category

1. Crime-Justice-Terrorism-Corruption
2. Freedom-Free speech-Resistance & H-rights
3. Health-Wellness-Healing-Nutrition & Fitness

- 4. Main
- 5. NWO-Deep State-Dictatorship-Tyrrany

Date Created
06/02/2022