



Quietly Published Pfizer Documents confirm between 82% & 97% of COVID Vaccinated Pregnant Women sadly lost their baby during the ongoing Clinical Trial

## Description

 Print  PDF  Email

The confidential Pfizer documents that the FDA have been forced to publish by court order reveal that 82% to 97% of women who were mistakenly exposed to the mRNA Covid-19 injection either suffered a miscarriage or suffered having to witness the death of their newborn child upon giving birth.

But Pfizer falsely claimed – “There were no safety signals that emerged from the review of these cases of use in pregnancy”.

BNT162b2

5.3.6 Cumulative Analysis of Post-authorization Adverse Event Reports

**Table 6. Description of Missing Information**

Topic	Description
Missing Information	<p><b>Post Authorization Cases Evaluation (cumulative to 28 Feb 2021)</b> <b>Total Number of Cases in the Reporting Period (N=42086)</b></p> <ul style="list-style-type: none"><li>In 4 cases (3 non-serious; 1 serious) Suppressed lactation occurred in a breast feeding women with the following co-reported events: Pyrexia (2), Paresis, Headache, Chills, Vomiting, Pain in extremity, Arthralgia, Breast pain, Scar pain, Nausea, Migraine, Myalgia, Fatigue and Breast milk discolouration (1 each).</li></ul> <p>Conclusion: There were no safety signals that emerged from the review of these cases of use in pregnancy and while breast feeding.</p>

The [US Food and Drug Administration](#) (FDA) attempted to delay the release of Pfizer’s COVID-19 vaccine safety data for 75 years despite approving the injection after only 108 days of safety review on [December 11th, 2020](#).

But in early January 2022, Federal Judge Mark Pittman ordered them to release 55,000 pages per

month. They released 12,000 pages by the end of January.

Since then, PHMPT has posted all of the [documents](#) on its website.

One of the documents contained in the data dump is '[reissue\\_5.3.6 postmarketing experience.pdf](#)'. Page 12 of the confidential document contains data on the use of the Pfizer Covid-19 injection in pregnancy and lactation.

Pfizer claim in the document that by 28th February 2021 there were 270 known cases of exposure to the mRNA injection during pregnancy.

**Table 6. Description of Missing Information**

Topic	Description
<b>Missing Information</b>	<b>Post Authorization Cases Evaluation (cumulative to 28 Feb 2021)</b> <b>Total Number of Cases in the Reporting Period (N=42086)</b>
Use in Pregnancy and lactation	<ul style="list-style-type: none"> <li>Number of cases: 413<sup>a</sup> (0.98% of the total PM dataset); 84 serious and 329 non-serious;</li> <li>Country of incidence: US (205), UK (64), Canada (31), Germany (30), Poland (13), Israel (11); Italy (9), Portugal (8), Mexico (6), Estonia, Hungary and Ireland, (5 each), Romania (4), Spain (3), Czech Republic and France (2 each), the remaining 10 cases were distributed among 10 other countries.</li> </ul> <p>Pregnancy cases: 274 cases including:</p> <ul style="list-style-type: none"> <li>270 mother cases and 4 fetus/baby cases representing 270 unique pregnancies (the 4 fetus/baby cases were linked to 3 mother cases; 1 mother case involved twins).</li> </ul>

One-hundred-and-forty-six of those mother cases did not immediately report the immediate occurrence of any clinical adverse event. But 124 of the 270 mother cases did. Meaning 46% of the mothers exposed to the Pfizer Covid-19 injection suffered an adverse reaction.

Of those 124 mothers suffering an adverse reaction, 49 were considered non-serious adverse reactions, whereas 75 were considered serious. This means 58% of the mothers who reported suffering adverse reactions suffered a serious adverse event ranging from uterine contraction to foetal death.

- 146 non-serious mother cases reported exposure to vaccine in utero without the occurrence of any clinical adverse event. The exposure PTs coded to the PTs Maternal exposure during pregnancy (111), Exposure during pregnancy (29) and Maternal exposure timing unspecified (6). Trimester of exposure was reported in 21 of these cases: 1st trimester (15 cases), 2nd trimester (7), and 3rd trimester (2).
- 124 mother cases, 49 non-serious and 75 serious, reported clinical events, which occurred in the vaccinated mothers. Pregnancy related events reported in these cases coded to the PTs Abortion spontaneous (25), Uterine contraction during pregnancy, Premature rupture of membranes, Abortion, Abortion missed, and Foetal death (1 each). Other clinical events which occurred in more than 5 cases coded to the PTs Headache (33), Vaccination site pain (24), Pain in extremity and Fatigue (22 each), Myalgia and Pyrexia (16 each), Chills (13) Nausea (12), Pain (11), Arthralgia (9), Lymphadenopathy and Drug ineffective (7 each), Chest pain, Dizziness and Asthenia (6 each), Malaise and COVID-19 (5 each). Trimester of exposure was reported in 22 of these cases: 1st trimester (19 cases), 2nd trimester (1 case), 3rd trimester (2 cases).

A total of 4 serious foetus/baby cases were reported due to exposure to the Pfizer injection. The 4 serious cases involved the following events –


- Foetal growth restriction x2
- Premature baby x2
- Neonatal Death x1
- 4 serious foetus/baby cases reported the PTs Exposure during pregnancy, Foetal growth restriction, Maternal exposure during pregnancy, Premature baby (2 each), and Death neonatal (1). Trimester of exposure was reported for 2 cases (twins) as occurring during the 1st trimester.

But here's where things get rather concerning. Pfizer state that of the 270 pregnancies they have absolutely no idea what happened in 238 of them.

each). No outcome was provided for 238 pregnancies (note that 2 different outcomes were reported for each twin, and both were counted).

But here are the known outcomes of the remaining pregnancies –

- Spontaneous Abortion (miscarriage) x23,
- Outcome pending x5,
- Premature baby with neonatal death x 2,
- Spontaneous Abortion with intrauterine death x2,
- Spontaneous Abortion with neonatal death x 1
- Normal outcome x1



- Pregnancy outcomes for the 270 pregnancies were reported as spontaneous abortion (23), outcome pending (5), premature birth with neonatal death, spontaneous abortion with intrauterine death (2 each), spontaneous abortion with neonatal death, and normal outcome (1 each). No outcome was provided for 238 pregnancies (note that 2 different outcomes were reported for each twin, and both were counted).

There were 34 outcomes altogether at the time of the report, but 5 of them were still pending. Pfizer note that only 1 of the 29 known outcomes were normal, whilst 28 of the 29 outcomes resulted in the loss/death of the baby. This equates to 97% of all known outcomes of Covid-19 vaccination during pregnancy resulting in the loss of the child.

When we include the 5 cases where the outcome was still pending it equates to 82% of all outcomes of Covid-19 vaccination during pregnancy resulting in the loss of the child. This equates to an average of around 90% between the 82% and 97% figure.

That 82% figure is, however, very interesting when we consider the outcome of another study conducted by the Centers for Disease Controls V-Safe COVID-19 Pregnancy Registry Team. A study that was used to justify not just offering, but harassing pregnant women to get the Covid-19 injection in the UK.

We [exclusively revealed](#) in July 2021 how data had been manipulated by scientists carrying out a [real world study](#)

for the CDC to show that Covid-19 vaccines were safe for use during pregnancy.

The authors claimed that the number of people to suffer a spontaneous abortion (miscarriage) during the study was 104 out of 827 completed pregnancies, equating the risk of miscarriage at 12.6%; 7 – 12% lower than the risk of miscarriage in the general population.

**Table 4**

**Pregnancy Loss and Neonatal Outcomes in Published Studies and V-safe Pregnancy Registry Participants.**

Participant-Reported Outcome	Published Incidence <sup>*</sup> %	V-safe Pregnancy Registry <sup>†</sup> no./total no. (%)
Pregnancy loss among participants with a completed pregnancy		
Spontaneous abortion: <20 wk <sup>15-17</sup>	10-26	104/827 (12.6) <sup>‡</sup>
Stillbirth: ≥ 20 wk <sup>18-20</sup>	<1	1/725 (0.1) <sup>§</sup>
Neonatal outcome among live-born infants		
Preterm birth: <37 wk <sup>21,22</sup>	8-15	60/636 (9.4) <sup>¶</sup>
Small size for gestational age <sup>23,24</sup>	3.5	23/724 (3.2)
Congenital anomalies <sup>25**</sup>	3	16/724 (2.2)
Neonatal death <sup>26††</sup>	<1	0/724

[Source](#)

However, our analysis proved that these numbers were extremely misleading due to the fact that of the 827 completed pregnancies, 700 / 86% of the women had received a dose of either the Pfizer or Moderna Covid-19 vaccine during the third trimester of pregnancy, meaning it was impossible for them to suffer a miscarriage due to the fact they can only occur prior to week 20 of pregnancy.

This meant that just 127 women received either the Pfizer or Moderna Covid-19 vaccine during the first/second trimester, with 104 of the woman sadly losing their baby.

Therefore [the rate of incidence of miscarriage was 82%](#), **not** 12.6% as presented in the findings of the study, and the authors of the study have since admitted that they made a mistake, [issuing a correction](#) six months too late because [the study](#) has been used to justify Covid-19 vaccination of pregnant women and new mothers around the world.



\*The populations from which these rates are derived are not matched to the current study population for age, race and ethnic group, or other demographic and clinical factors.

†Data on pregnancy loss are based on 827 participants in the v-safe pregnancy registry who received an mRNA Covid-19 vaccine (BNT162b2 [Pfizer–BioNTech] or mRNA-1273 [Moderna]) from December 14, 2020, to February 28, 2021, and who reported a completed pregnancy. A total of 700 participants (84.6%) received their first eligible dose in the third trimester. Data on neonatal outcomes are based on 724 live-born infants, including 12 sets of multiples.

‡A total of 96 of 104 spontaneous abortions (92.3%) occurred before 13 weeks of gestation.

§The denominator includes live-born infants and stillbirths.

¶The denominator includes only participants vaccinated before 37 weeks of gestation.

‡Small size for gestational age indicates a birthweight below the 10th percentile for gestational age and infant sex according to INTERGROWTH-21<sup>st</sup> growth standards (<http://intergrowth21.ndog.ox.ac.uk>). These standards draw from an international sample including both low-income and high-income countries but exclude children with coexisting conditions and malnutrition. They can be used as a standard for healthy children growing under optimal conditions.

\*\*Values include only major congenital anomalies in accordance with the Metropolitan Atlanta Congenital Defects Program 6-Digit Code Defect List ([www.cdc.gov/ncbddd/birthdefects/macdp.html](http://www.cdc.gov/ncbddd/birthdefects/macdp.html)); all pregnancies with major congenital anomalies were exposed to Covid-19 vaccines only in the third trimester of pregnancy (i.e., well after the period of organogenesis).

††Neonatal death indicates death within the first 28 days after delivery.

### Source

But there's that 82% number again. So that's two sets of data, Pfizer's confidential data, and the CDC's manipulated study that show the rate of pregnancy loss following Covid-19 vaccination to be 82%. Just a coincidence?

Highly unlikely when we consider what else Medicine Regulators and Pfizer tried to cover up.

A 'Freedom of Information' request alongside an in-depth dive into the only pregnancy/fertility study performed on the Pfizer Covid-19 injection reveals that Pfizer and Medicine Regulators hid the dangers of Covid-19 Vaccination during Pregnancy because the study found it increases the risk of birth defects and infertility.

You can read the full investigation [here](#), but here's a snippet of what should have been included in official public documents, and subsequently wasn't –

As discussed in the assessment, Pregnancy Category B2 is considered appropriate for this product. Following changes are recommended.

“Pregnancy Category B2

There is limited experience with use of COMIRNATY in pregnant women. ~~Animal studies do~~ A combined fertility and developmental toxicity study in rats showed increased occurrence of supernumerary lumbar ribs in fetuses from COMIRNATY-treated female rats did not indicate direct or indirect harmful effects with respect to pregnancy, embryo/fetal development, r post-natal development (see Effects on fertility Section 5.3 Preclinical safety data). Administration of COMIRNATY in pregnancy should only be considered when the potential benefits outweigh any potential risks for the mother and fetus.”

[Source](#)




The FDA’S decision not to act on Pfizer’s data, and the cover-up of the animal study findings by both Pfizer and Medicine Regulators around the world has led to [4,113 foetal deaths being reported in the USA alone](#) as adverse reactions to the Covid-19 injections as of 22nd April 2022.

# 4,113 Fetal Deaths

## *following COVID-19 vaccines*


### 17 Months Time Period

Table

		
Vaccine/Manufacturer	Count	Percent
<b>TOTAL</b>	† 4,382	† 106.54%
COVID19 / PFIZER/BIONTECH	3,209	78.02%
COVID19 / MODERNA	996	24.22%
COVID19 / JANSSEN	140	3.4%
UNK / UNKNOWN MANUFACTURER	21	0.51%
COVID19 / UNKNOWN MANUFACTURER	10	0.24%
HEPA / UNKNOWN MANUFACTURER	2	0.05%
TDAP / GLAXOSMITHKLINE BIOLOGICALS	1	0.02%
TD / SANOFI PASTEUR	1	0.02%
IPV / SANOFI PASTEUR	1	0.02%
FLUC4 / SEQIRUS, INC.	1	0.02%
† Because some cases have multiple vaccinations and symptoms, a single case can account for multiple entries in this table. This is the reason why the Total Count is greater than 4113 (the number of cases found), and the Total Percentage is greater than 100.		

And take a look at this [data coming out of Scotland](#) –





*Click to enlarge*

Public Health Scotland (PHS) has a full dashboard on Covid-19 wider impacts on the health care system, found [here](#), and it includes a whole range of data from mental health statistics to pregnancies, cardiovascular disorders data, and cancer.

[Official figures](#) reveal that the rate of neonatal deaths increased to 4.6 per 1000 live births in March 2022, a 119% increase on the expected rate of deaths. This means the neonatal mortality rate breached an upper warning threshold known as the 'control limit' for the second time in at least four years.

The last time it breached was in September 2021, when neonatal deaths per 1000 live births climbed to 5.1. Although the rate fluctuates month to month, the figure for both September 2021 and March 2022 is on a par with levels that were last typically seen in the late 1980s.

Public Health Scotland (PHS) has not formally announced they have launched an investigation, but this is what they are supposed to do when the upper warning threshold is reached, and they did so back in 2021.

At the time, PHS said the fact that the upper control limit has been exceeded "indicates there is a higher likelihood that there are factors beyond random variation that may have contributed to the number of deaths that occurred". You can read more [here](#).

We either have a serious issue here or a bucket load of terrible coincidences, and it's all arisen thanks to this unbelievable claim made by Pfizer in their confidential documents in response to 82% to 97% of women mistakenly exposed to the Covid-19 injection during pregnancy losing their child –

"There were no safety signals that emerged from the review of these cases of use in pregnancy and while breast feeding."

BNT162b2

5.3.6 Cumulative Analysis of Post-authorization Adverse Event Reports

**Table 6. Description of Missing Information**

Topic	Description
Missing Information	<b>Post Authorization Cases Evaluation (cumulative to 28 Feb 2021)</b> <b>Total Number of Cases in the Reporting Period (N=42086)</b>
	<ul style="list-style-type: none"><li>In 4 cases (3 non-serious; 1 serious) Suppressed lactation occurred in a breast feeding women with the following co-reported events: Pyrexia (2), Paresis, Headache, Chills, Vomiting, Pain in extremity, Arthralgia, Breast pain, Scar pain, Nausea, Migraine, Myalgia, Fatigue and Breast milk discolouration (1 each).</li></ul> <p>Conclusion: There were no safety signals that emerged from the review of these cases of use in pregnancy and while breast feeding.</p>

See for yourself on pages 12 and 13 of the confidential document found [here](#).

**Category**

1. Crime-Justice-Terrorism-Corruption
2. Disasters-Crisis-Depopulation-Genocide
3. Health-Wellness-Healing-Nutrition & Fitness
4. Main
5. NWO-Deep State-Dictatorship-Tyranny

**Date Created**

04/16/2023